

# Refining Jensen's Integral Inequality for Partitions of Weights

Silvestru Sever Dragomir<sup>1,2</sup>

**Abstract:** In this paper we establish a refinement and some reverses for Jensen's inequality for the general Lebesgue integral on measurable spaces and partitions of weights. Applications for discrete inequalities and weighted means of positive numbers are also given.

**Keywords:** Jensen's inequality, Convex functions, Lebesgue integral, Weighted means.

**MSC2010:** Primary 26D15; Secondary 94A17.

## 1 Introduction

Let  $C$  be a convex subset of the linear space  $X$  and  $f$  a convex function on  $C$ . If  $\mathbf{p} = (p_1, \dots, p_n)$  is a *probability sequence*, i.e.  $p_1, \dots, p_n \geq 0$  with  $\sum_{i=1}^n p_i = 1$  and  $\mathbf{x} = (x_1, \dots, x_n) \in C^n$ , then

$$f\left(\sum_{i=1}^n p_i x_i\right) \leq \sum_{i=1}^n p_i f(x_i), \quad (1.1)$$

is well known in the literature as *Jensen's inequality*.

The Jensen inequality for convex functions plays a crucial role in the Theory of Inequalities due to the fact that other inequalities such as the arithmetic mean-geometric mean inequality, Hölder and Minkowski's inequalities, Ky Fan's inequality etc. can be obtained as particular cases of it.

In 1989, J. Pečarić and the author obtained the following refinement of (1.1) (see [14]):

$$\begin{aligned} f\left(\sum_{i=1}^n p_i x_i\right) &\leq \sum_{i_1, \dots, i_{k+1}=1}^n p_{i_1} \dots p_{i_{k+1}} f\left(\frac{x_{i_1} + \dots + x_{i_{k+1}}}{k+1}\right) \\ &\leq \sum_{i_1, \dots, i_k=1}^n p_{i_1} \dots p_{i_k} f\left(\frac{x_{i_1} + \dots + x_{i_k}}{k}\right) \leq \dots \leq \sum_{i=1}^n p_i f(x_i), \end{aligned} \quad (1.2)$$

for  $k \geq 1$  and  $\mathbf{p}, \mathbf{x}$  as above.

If  $q_1, \dots, q_k \geq 0$  with  $\sum_{j=1}^k q_j = 1$ , then the following weighted refinement obtained in 1994 by the author also holds (see [5]):

$$\begin{aligned} f\left(\sum_{i=1}^n p_i x_i\right) &\leq \sum_{i_1, \dots, i_k=1}^n p_{i_1} \dots p_{i_k} f\left(\frac{x_{i_1} + \dots + x_{i_k}}{k}\right) \\ &\leq \sum_{i_1, \dots, i_k=1}^n p_{i_1} \dots p_{i_k} f(q_1 x_{i_1} + \dots + q_k x_{i_k}) \leq \sum_{i=1}^n p_i f(x_i), \end{aligned} \quad (1.3)$$

where  $1 \leq k \leq n$  and  $\mathbf{p}, \mathbf{x}$  are as above.

More recently the author obtained a different refinement of Jensen's inequality incorporated in (see [7]):

$$\begin{aligned} f\left(\sum_{j=1}^n p_j x_j\right) &\leq \min_{k \in \{1, \dots, n\}} \left[ (1 - p_k) f\left(\frac{\sum_{j=1}^n p_j x_j - p_k x_k}{1 - p_k}\right) + p_k f(x_k) \right] \\ &\leq \frac{1}{n} \left[ \sum_{k=1}^n (1 - p_k) f\left(\frac{\sum_{j=1}^n p_j x_j - p_k x_k}{1 - p_k}\right) + \sum_{k=1}^n p_k f(x_k) \right] \\ &\leq \max_{k \in \{1, \dots, n\}} \left[ (1 - p_k) f\left(\frac{\sum_{j=1}^n p_j x_j - p_k x_k}{1 - p_k}\right) + p_k f(x_k) \right] \\ &\leq \sum_{j=1}^n p_j f(x_j), \end{aligned} \quad (1.4)$$

where  $f, x_k$  and  $p_k$  are as above.

For other refinements and applications related to Ky Fan's inequality, the arithmetic mean-geometric mean inequality, the generalized triangle inequality, the  $f$ -Divergence measure etc., see [1], [2]-[12], [13] and [15]-[16]

Motivated by the above results, we investigate in this paper the integral version of Jensen inequality and establish some refinements and reverses of interest for applications.

Let  $(\Omega, \mathcal{A}, \mu)$  be a measurable space consisting of a set  $\Omega$ , a  $\sigma$ -algebra  $\mathcal{A}$  of parts of  $\Omega$  and a countably additive and positive measure  $\mu$  on  $\mathcal{A}$  with values in  $\mathbb{R} \cup \{\infty\}$ . For the  $\mu$ -integrable positive  $\mu$ -a.e. weight  $w$  consider the Lebesgue space

$$L_w(\Omega, \mu) := \{f : \Omega \rightarrow \mathbb{R}, f \text{ is } \mu\text{-measurable and } \int_{\Omega} |f(t)| w(t) d\mu(t) < \infty\}.$$

For simplicity of notation we write everywhere in the sequel  $\int_{\Omega} w d\mu$  instead of  $\int_{\Omega} w(t) d\mu(t)$  etc..

For the  $\mu$ -integrable positive  $\mu$ -a.e. weight  $w$  and a given  $n \geq 2$  we consider the set  $\mathfrak{P}_n(w)$  all possible  $n$ -tuples of  $\mu$ -integrable positive  $\mu$ -a.e. weights  $\bar{w} = (w_1, \dots, w_n)$  with the property that  $\sum_{i=1}^n w_i = w$ . The  $n$ -tuple  $\bar{w} = (w_1, \dots, w_n)$  it is called a *partition* of the weight  $w$ . It is clear that  $\sum_{i=1}^n \int_{\Omega} w_i d\mu = \int_{\Omega} w d\mu$  for any  $(w_1, \dots, w_n) \in \mathfrak{P}_n(w)$  and  $\int_{\Omega} w_i d\mu > 0$ .

For a convex function  $\Phi : [m, M] \rightarrow \mathbb{R}$ , a  $\mu$ -measurable function  $f : \Omega \rightarrow [m, M]$  such that  $f, \Phi \circ f \in L_w(\Omega, \mu)$  we define the functional  $\psi(\Phi, f, \cdot) : \mathfrak{P}_n(w) \rightarrow \mathbb{R}$  by

$$\psi(\Phi, f, \bar{w}) := \frac{1}{\int_{\Omega} w d\mu} \sum_{i=1}^n \Phi \left( \frac{\int_{\Omega} f w_i d\mu}{\int_{\Omega} w_i d\mu} \right) \int_{\Omega} w_i d\mu. \quad (1.5)$$

In the next section we establish some results concerning this functional that are related to Jensen's integral inequality. Applications for discrete inequalities and weighted means are provided in the third section. In the last section some applications for univariate functions are also given.

## 2 Main Results

The following basic result holds:

**Theorem 2.1.** *Let  $\Phi : [m, M] \rightarrow \mathbb{R}$  be a convex function,  $f : \Omega \rightarrow [m, M]$  a  $\mu$ -measurable function such that  $f, \Phi \circ f \in L_w(\Omega, \mu)$ . Then for any  $\bar{w} \in \mathfrak{P}_n(w)$  we have*

$$\frac{\int_{\Omega} (\Phi \circ f) w d\mu}{\int_{\Omega} w d\mu} \geq \psi(\Phi, f, \bar{w}) \geq \Phi \left( \frac{\int_{\Omega} f w d\mu}{\int_{\Omega} w d\mu} \right), \quad (2.1)$$

where  $n \geq 2$ .

*Proof.* From Jensen's integral inequality we have

$$\int_{\Omega} (\Phi \circ f) w_i d\mu \geq \Phi \left( \frac{\int_{\Omega} f w_i d\mu}{\int_{\Omega} w_i d\mu} \right) \int_{\Omega} w_i d\mu \quad (2.2)$$

for any  $i \in \{1, \dots, n\}$ .

If we sum the inequality (2.2) over  $i$  from 1 to  $n$  we get

$$\sum_{i=1}^n \int_{\Omega} (\Phi \circ f) w_i d\mu \geq \sum_{i=1}^n \Phi \left( \frac{\int_{\Omega} f w_i d\mu}{\int_{\Omega} w_i d\mu} \right) \int_{\Omega} w_i d\mu \quad (2.3)$$

and since

$$\sum_{i=1}^n \int_{\Omega} (\Phi \circ f) w_i d\mu = \int_{\Omega} (\Phi \circ f) \left( \sum_{i=1}^n w_i \right) d\mu = \int_{\Omega} (\Phi \circ f) w d\mu$$

then from (2.3) we get the first part of (2.1).

Let

$$p_i = \int_{\Omega} w_i d\mu > 0 \quad z_i = \frac{\int_{\Omega} f w_i d\mu}{\int_{\Omega} w_i d\mu} \in [m, M], \quad i \in \{1, \dots, n\}.$$

Then

$$P_n := \sum_{i=1}^n p_i = \int_{\Omega} w d\mu,$$

and

$$\sum_{i=1}^n p_i z_i = \sum_{i=1}^n \int_{\Omega} f w_i d\mu = \int_{\Omega} f w d\mu.$$

From Jensen's discrete inequality

$$\frac{1}{P_n} \sum_{i=1}^n p_i \Phi(z_i) \geq \Phi\left(\frac{\sum_{i=1}^n p_i z_i}{P_n}\right)$$

we have

$$\frac{1}{\int_{\Omega} w d\mu} \sum_{i=1}^n \Phi\left(\frac{\int_{\Omega} f w_i d\mu}{\int_{\Omega} w_i d\mu}\right) \int_{\Omega} w_i d\mu \geq \Phi\left(\frac{\int_{\Omega} f w d\mu}{\int_{\Omega} w d\mu}\right)$$

and the second inequality in (2.1) is also proved.  $\square$

**Remark 2.2.** The double inequality (2.1) is equivalent to

$$\frac{\int_{\Omega} (\Phi \circ f) w d\mu}{\int_{\Omega} w d\mu} \geq \sup_{\bar{w} \in \mathfrak{P}_n(w)} \psi(\Phi, f, \bar{w}) \quad (2.4)$$

and

$$\inf_{\bar{w} \in \mathfrak{P}_n(w)} \psi(\Phi, f, \bar{w}) \geq \Phi\left(\frac{\int_{\Omega} f w d\mu}{\int_{\Omega} w d\mu}\right), \quad (2.5)$$

where  $n \geq 2$ .

We use the following lemma [9].

**Lemma 2.3.** *Let  $\Phi : I \rightarrow \mathbb{R}$  be a convex function on the interval of real numbers  $I$  and  $m, M \in \mathbb{R}$ ,  $m < M$  with  $[m, M] \subset I$ ,  $I$  is the interior of  $I$ . If  $g : \Omega \rightarrow \mathbb{R}$  is  $\mu$ -measurable, satisfies the bounds*

$$-\infty < m \leq g(x) \leq M < \infty \text{ for } \mu\text{-a.e. } x \in \Omega$$

and such that  $g, \Phi \circ g \in L_p(\Omega, \mu)$ , where  $p \geq 0$   $\mu$ -a.e. on  $\Omega$  with  $\int_{\Omega} p d\mu = 1$ , then

$$\begin{aligned}
0 &\leq \int_{\Omega} p (\Phi \circ f) d\mu - \Phi \left( \int_{\Omega} p f d\mu \right) \\
&\leq \frac{(M - \int_{\Omega} p f d\mu) (\int_{\Omega} p f d\mu - m)}{M - m} \sup_{t \in (m, M)} \Psi_{\Phi}(t; m, M) \\
&\leq \left( M - \int_{\Omega} p f d\mu \right) \left( \int_{\Omega} p f d\mu - m \right) \frac{\Phi'_-(M) - \Phi'_+(m)}{M - m} \\
&\leq \frac{1}{4} (M - m) [\Phi'_-(M) - \Phi'_+(m)],
\end{aligned} \tag{2.6}$$

where  $\Psi_{\Phi}(\cdot; m, M) : (m, M) \rightarrow \mathbb{R}$  is defined by

$$\Psi_{\Phi}(t; m, M) = \frac{\Phi(M) - \Phi(t)}{M - t} - \frac{\Phi(t) - \Phi(m)}{t - m}.$$

We have the following reverse of the first inequality in (2.1).

**Theorem 2.4.** *Let  $\Phi : [m, M] \rightarrow \mathbb{R}$  be a convex function,  $f : \Omega \rightarrow [m, M]$  a  $\mu$ -measurable function such that  $f, \Phi \circ f \in L_w(\Omega, \mu)$ . Then for any  $\bar{w} \in \mathfrak{P}_n(w)$ ,  $n \geq 2$  we have*

$$\begin{aligned}
0 &\leq \frac{\int_{\Omega} (\Phi \circ f) w d\mu}{\int_{\Omega} w d\mu} - \psi(\Phi, f, \bar{w}) \\
&\leq \frac{1}{M - m} \sup_{t \in (m, M)} \Psi_{\Phi}(t; m, M) \\
&\quad \times \frac{1}{\int_{\Omega} w d\mu} \sum_{i=1}^n \left( \int_{\Omega} w_i d\mu \right) \left( M - \frac{\int_{\Omega} w_i f d\mu}{\int_{\Omega} w_i d\mu} \right) \left( \frac{\int_{\Omega} w_i f d\mu}{\int_{\Omega} w_i d\mu} - m \right) \\
&\leq \frac{1}{M - m} \sup_{t \in (m, M)} \Psi_{\Phi}(t; m, M) \left( M - \frac{\int_{\Omega} f w d\mu}{\int_{\Omega} w d\mu} \right) \left( \frac{\int_{\Omega} f w d\mu}{\int_{\Omega} w d\mu} - m \right).
\end{aligned} \tag{2.7}$$

*Proof.* From the second inequality in (2.6) for  $g = f$  and  $p = \frac{w_i}{\int_{\Omega} w_i d\mu}$ ,  $i \in \{1, \dots, n\}$  we have

$$\begin{aligned}
0 &\leq \frac{1}{\int_{\Omega} w_i d\mu} \int_{\Omega} w_i (\Phi \circ f) d\mu - \Phi \left( \frac{1}{\int_{\Omega} w_i d\mu} \int_{\Omega} w_i f d\mu \right) \\
&\leq \frac{1}{M - m} \left( M - \frac{\int_{\Omega} w_i f d\mu}{\int_{\Omega} w_i d\mu} \right) \left( \frac{\int_{\Omega} w_i f d\mu}{\int_{\Omega} w_i d\mu} - m \right) \sup_{t \in (m, M)} \Psi_{\Phi}(t; m, M)
\end{aligned} \tag{2.8}$$

for any  $i \in \{1, \dots, n\}$ .

If we multiply by  $\int_{\Omega} w_i d\mu > 0$  and sum over  $i$  from 1 to  $n$  we get

$$\begin{aligned}
0 &\leq \sum_{i=1}^n \int_{\Omega} w_i (\Phi \circ f) d\mu - \sum_{i=1}^n \Phi \left( \frac{1}{\int_{\Omega} w_i d\mu} \int_{\Omega} w_i f d\mu \right) \int_{\Omega} w_i d\mu & (2.9) \\
&\leq \frac{1}{M-m} \sup_{t \in (m, M)} \Psi_{\Phi}(t; m, M) \\
&\quad \times \sum_{i=1}^n \left( \int_{\Omega} w_i d\mu \right) \left( M - \frac{\int_{\Omega} w_i f d\mu}{\int_{\Omega} w_i d\mu} \right) \left( \frac{\int_{\Omega} w_i f d\mu}{\int_{\Omega} w_i d\mu} - m \right),
\end{aligned}$$

which proves the second inequality in (2.7).

Now, observe that the function  $\Psi : [m, M] \rightarrow [0, \infty)$ ,  $\Psi(t) = (M-t)(t-m)$  is concave and by Jensen's inequality for concave functions with

$$p_i = \int_{\Omega} w_i d\mu > 0 \quad z_i = \frac{\int_{\Omega} f w_i d\mu}{\int_{\Omega} w_i d\mu} \in [m, M], \quad i \in \{1, \dots, n\}$$

we have

$$\begin{aligned}
&\frac{1}{\int_{\Omega} w d\mu} \sum_{i=1}^n \left( \int_{\Omega} w_i d\mu \right) \left( M - \frac{\int_{\Omega} w_i f d\mu}{\int_{\Omega} w_i d\mu} \right) \left( \frac{\int_{\Omega} w_i f d\mu}{\int_{\Omega} w_i d\mu} - m \right) \\
&\leq \left( M - \frac{1}{\int_{\Omega} w d\mu} \sum_{i=1}^n \left( \int_{\Omega} w_i d\mu \right) \frac{\int_{\Omega} w_i f d\mu}{\int_{\Omega} w_i d\mu} \right) \\
&\quad \times \left( \frac{1}{\int_{\Omega} w d\mu} \sum_{i=1}^n \left( \int_{\Omega} w_i d\mu \right) \frac{\int_{\Omega} w_i f d\mu}{\int_{\Omega} w_i d\mu} - m \right) \\
&= \left( M - \frac{\int_{\Omega} f w d\mu}{\int_{\Omega} w d\mu} \right) \left( \frac{\int_{\Omega} f w d\mu}{\int_{\Omega} w d\mu} - m \right),
\end{aligned}$$

which proves the last part of (2.7). □

**Remark 2.5.** Since, as shown in [9],

$$\sup_{t \in (m, M)} \Psi_{\Phi}(t; m, M) \leq \Phi'_-(M) - \Phi'_+(m)$$

then we have the following simpler inequality

$$\begin{aligned}
0 &\leq \frac{\int_{\Omega} (\Phi \circ f) w d\mu}{\int_{\Omega} w d\mu} - \psi(\Phi, f, \bar{w}) \\
&\leq \frac{\Phi'_-(M) - \Phi'_+(m)}{M - m} \\
&\quad \times \sum_{i=1}^n \frac{\int_{\Omega} w_i d\mu}{\int_{\Omega} w d\mu} \left( M - \frac{\int_{\Omega} w_i f d\mu}{\int_{\Omega} w_i d\mu} \right) \left( \frac{\int_{\Omega} w_i f d\mu}{\int_{\Omega} w_i d\mu} - m \right) \\
&\leq \frac{\Phi'_-(M) - \Phi'_+(m)}{M - m} \left( M - \frac{\int_{\Omega} f w d\mu}{\int_{\Omega} w d\mu} \right) \left( \frac{\int_{\Omega} f w d\mu}{\int_{\Omega} w d\mu} - m \right).
\end{aligned} \tag{2.10}$$

If we use Lemma 2.3 for the discrete measure, we can state the following result:

**Lemma 2.6.** *Let  $\Phi : [a, b] \rightarrow \mathbb{R}$  be a convex function on the interval of real numbers  $[a, b]$ ,  $z_i \in [a, b]$ ,  $p_i \geq 0$ ,  $i \in \{1, \dots, n\}$  and  $\sum_{i=1}^n p_i = 1$ . Then*

$$\begin{aligned}
0 &\leq \sum_{i=1}^n p_i \Phi(z_i) - \Phi \left( \sum_{i=1}^n p_i z_i \right) \\
&\leq \frac{(b - \sum_{i=1}^n p_i z_i) (\sum_{i=1}^n p_i z_i - a)}{b - a} \sup_{t \in (a, b)} \Psi_{\Phi}(t; a, b) \\
&\leq \left( b - \sum_{i=1}^n p_i z_i \right) \left( \sum_{i=1}^n p_i z_i - a \right) \frac{\Phi'_-(b) - \Phi'_+(a)}{b - a} \\
&\leq \frac{1}{4} (b - a) [\Phi'_-(b) - \Phi'_+(a)].
\end{aligned} \tag{2.11}$$

The following reverse of the second inequality in (2.1) holds:

**Theorem 2.7.** *Let  $\Phi : [m, M] \rightarrow \mathbb{R}$  be a convex function,  $f : \Omega \rightarrow [m, M]$  a  $\mu$ -measurable function such that  $f, \Phi \circ f \in L_w(\Omega, \mu)$ . Then for any  $\bar{w} \in \mathfrak{B}_n(w)$ ,  $n \geq 2$  we have*

$$\begin{aligned}
0 &\leq \psi(\Phi, f, \bar{w}) - \Phi \left( \frac{\int_{\Omega} f w d\mu}{\int_{\Omega} w d\mu} \right) \\
&\leq \frac{\left( L(\bar{w}) - \frac{\int_{\Omega} f w d\mu}{\int_{\Omega} w d\mu} \right) \left( \frac{\int_{\Omega} f w d\mu}{\int_{\Omega} w d\mu} - l(\bar{w}) \right)}{L(\bar{w}) - l(\bar{w})} \sup_{t \in (l(\bar{w}), L(\bar{w}))} \Psi_{\Phi}(t; l(\bar{w}), L(\bar{w})) \\
&\leq \frac{1}{4} (L(\bar{w}) - l(\bar{w})) \sup_{t \in (l(\bar{w}), L(\bar{w}))} \Psi_{\Phi}(t; l(\bar{w}), L(\bar{w})),
\end{aligned} \tag{2.12}$$

where

$$l(\bar{w}) := \min_{i \in \{1, \dots, n\}} \left\{ \frac{\int_{\Omega} f w_i d\mu}{\int_{\Omega} w_i d\mu} \right\}, \quad L(\bar{w}) := \max_{i \in \{1, \dots, n\}} \left\{ \frac{\int_{\Omega} f w_i d\mu}{\int_{\Omega} w_i d\mu} \right\}. \quad (2.13)$$

*Proof.* If we write the first two inequalities in (2.11) for

$$p_i = \frac{\int_{\Omega} w_i d\mu}{\int_{\Omega} w d\mu} > 0 \quad z_i = \frac{\int_{\Omega} f w_i d\mu}{\int_{\Omega} w_i d\mu}, \quad i \in \{1, \dots, n\}$$

and for  $a = l(\bar{w})$ ,  $b = L(\bar{w})$  as above we have

$$\begin{aligned} 0 &\leq \frac{1}{\int_{\Omega} w d\mu} \sum_{i=1}^n \left( \int_{\Omega} w_i d\mu \right) \Phi \left( \frac{\int_{\Omega} f w_i d\mu}{\int_{\Omega} w_i d\mu} \right) - \Phi \left( \frac{\sum_{i=1}^n \int_{\Omega} f w_i d\mu}{\int_{\Omega} w d\mu} \right) \\ &\leq \frac{\sup_{t \in (l(\bar{w}), L(\bar{w}))} \Psi_{\Phi}(t; l(\bar{w}), L(\bar{w}))}{L(\bar{w}) - l(\bar{w})} \\ &\quad \times \left( L(\bar{w}) - \sum_{i=1}^n \frac{\int_{\Omega} w_i d\mu}{\int_{\Omega} w d\mu} \cdot \frac{\int_{\Omega} f w_i d\mu}{\int_{\Omega} w_i d\mu} \right) \left( \sum_{i=1}^n \frac{\int_{\Omega} w_i d\mu}{\int_{\Omega} w d\mu} \cdot \frac{\int_{\Omega} f w_i d\mu}{\int_{\Omega} w_i d\mu} - l(\bar{w}) \right) \\ &= \frac{\left( L(\bar{w}) - \frac{\int_{\Omega} f w d\mu}{\int_{\Omega} w d\mu} \right) \left( \frac{\int_{\Omega} f w d\mu}{\int_{\Omega} w d\mu} - l(\bar{w}) \right)}{L(\bar{w}) - l(\bar{w})} \sup_{t \in (l(\bar{w}), L(\bar{w}))} \Psi_{\Phi}(t; l(\bar{w}), L(\bar{w})), \end{aligned} \quad (2.14)$$

which proves the second inequality in (2.12).

The last part in (2.12) follows by the elementary inequality

$$\alpha\beta \leq \left( \frac{\alpha + \beta}{2} \right)^2, \quad \alpha, \beta \in \mathbb{R}.$$

□

**Remark 2.8.** Since

$$\Psi_{\Phi}(t; l(\bar{w}), L(\bar{w})) \leq \Phi'_{-}(L(\bar{w})) - \Phi'_{+}(l(\bar{w})),$$

then from (2.12) we have the simpler inequalities

$$\begin{aligned} 0 &\leq \psi(\Phi, f, \bar{w}) - \Phi \left( \frac{\int_{\Omega} f w d\mu}{\int_{\Omega} w d\mu} \right) \\ &\leq \frac{\Phi'_{-}(L(\bar{w})) - \Phi'_{+}(l(\bar{w}))}{L(\bar{w}) - l(\bar{w})} \left( L(\bar{w}) - \frac{\int_{\Omega} f w d\mu}{\int_{\Omega} w d\mu} \right) \left( \frac{\int_{\Omega} f w d\mu}{\int_{\Omega} w d\mu} - l(\bar{w}) \right) \\ &\leq \frac{1}{4} (L(\bar{w}) - l(\bar{w})) [\Phi'_{-}(L(\bar{w})) - \Phi'_{+}(l(\bar{w}))]. \end{aligned} \quad (2.15)$$

The following reverse of Jensen inequality also holds [9]:

**Lemma 2.9.** *Let  $\Phi : I \rightarrow \mathbb{R}$  be a convex function on the interval of real numbers  $I$  and  $m, M \in \mathbb{R}$ ,  $m < M$  with  $[m, M] \subset \overset{\circ}{I}$ ,  $\overset{\circ}{I}$  is the interior of  $I$ . If  $g : \Omega \rightarrow \mathbb{R}$  is  $\mu$ -measurable, satisfies the bounds*

$$-\infty < m \leq g(x) \leq M < \infty \text{ for } \mu\text{-a.e. } x \in \Omega$$

and such that  $g, \Phi \circ g \in L_p(\Omega, \mu)$ , where  $p \geq 0$   $\mu$ -a.e. on  $\Omega$  with  $\int_{\Omega} p d\mu = 1$ , then

$$\begin{aligned} 0 &\leq \int_{\Omega} p(\Phi \circ f) d\mu - \Phi \left( \int_{\Omega} p f d\mu \right) & (2.16) \\ &\leq \left[ \frac{\Phi(m) + \Phi(M)}{2} - \Phi \left( \frac{m+M}{2} \right) \right] \left( 1 + \frac{2}{M-m} \left| \int_{\Omega} p f d\mu - \frac{m+M}{2} \right| \right) \\ &\leq 2 \left[ \frac{\Phi(m) + \Phi(M)}{2} - \Phi \left( \frac{m+M}{2} \right) \right]. \end{aligned}$$

We have the following result:

**Theorem 2.10.** *Let  $\Phi : [m, M] \rightarrow \mathbb{R}$  be a convex function,  $f : \Omega \rightarrow [m, M]$  a  $\mu$ -measurable function such that  $f, \Phi \circ f \in L_w(\Omega, \mu)$ . Then for any  $\bar{w} \in \mathfrak{P}_n(w)$ ,  $n \geq 2$  we have*

$$\begin{aligned} 0 &\leq \frac{\int_{\Omega} (\Phi \circ f) w d\mu}{\int_{\Omega} w d\mu} - \psi(\Phi, f, \bar{w}) & (2.17) \\ &\leq \left[ \frac{\Phi(m) + \Phi(M)}{2} - \Phi \left( \frac{m+M}{2} \right) \right] \\ &\quad \times \left( 1 + \frac{2}{M-m} \sum_{i=1}^n \left| \frac{1}{\int_{\Omega} w d\mu} \int_{\Omega} w_i \left( f - \frac{m+M}{2} \right) d\mu \right| \right) \\ &\leq \left[ \frac{\Phi(m) + \Phi(M)}{2} - \Phi \left( \frac{m+M}{2} \right) \right] \\ &\quad \times \left( 1 + \frac{2}{M-m} \frac{1}{\int_{\Omega} w d\mu} \int_{\Omega} w \left| f - \frac{m+M}{2} \right| d\mu \right). \end{aligned}$$

*Proof.* From the second inequality in (2.16) for  $g = f$  and  $p = \frac{w_i}{\int_{\Omega} w_i d\mu}$ ,  $i \in \{1, \dots, n\}$

we have

$$\begin{aligned}
0 &\leq \frac{1}{\int_{\Omega} w_i d\mu} \int_{\Omega} w_i (\Phi \circ f) d\mu - \Phi \left( \frac{1}{\int_{\Omega} w_i d\mu} \int_{\Omega} w_i f d\mu \right) \\
&\leq \left[ \frac{\Phi(m) + \Phi(M)}{2} - \Phi \left( \frac{m+M}{2} \right) \right] \\
&\quad \times \left( 1 + \frac{2}{M-m} \left| \frac{1}{\int_{\Omega} w_i d\mu} \int_{\Omega} w_i f d\mu - \frac{m+M}{2} \right| \right)
\end{aligned} \tag{2.18}$$

for any  $i \in \{1, \dots, n\}$ .

If we multiply the inequality (2.18) by  $\int_{\Omega} w_i d\mu > 0$  and sum over  $i$  from 1 to  $n$  we get

$$\begin{aligned}
0 &\leq \sum_{i=1}^n \int_{\Omega} w_i (\Phi \circ f) d\mu - \sum_{i=1}^n \Phi \left( \frac{1}{\int_{\Omega} w_i d\mu} \int_{\Omega} w_i f d\mu \right) \int_{\Omega} w_i d\mu \\
&\leq \left[ \frac{\Phi(m) + \Phi(M)}{2} - \Phi \left( \frac{m+M}{2} \right) \right] \\
&\quad \times \left( \sum_{i=1}^n \int_{\Omega} w_i d\mu + \frac{2}{M-m} \sum_{i=1}^n \left( \int_{\Omega} w_i d\mu \right) \left| \frac{1}{\int_{\Omega} w_i d\mu} \int_{\Omega} w_i f d\mu - \frac{m+M}{2} \right| \right) \\
&= \left[ \frac{\Phi(m) + \Phi(M)}{2} - \Phi \left( \frac{m+M}{2} \right) \right] \\
&\quad \times \left( \int_{\Omega} w d\mu + \frac{2}{M-m} \sum_{i=1}^n \left| \int_{\Omega} w_i \left( f - \frac{m+M}{2} \right) d\mu \right| \right)
\end{aligned}$$

any by dividing with  $\int_{\Omega} w d\mu = \sum_{i=1}^n \int_{\Omega} w_i d\mu$ , we get the second inequality in (2.17).

By the properties of modulus we have

$$\begin{aligned}
\sum_{i=1}^n \left| \frac{1}{\int_{\Omega} w d\mu} \int_{\Omega} w_i \left( f - \frac{m+M}{2} \right) d\mu \right| &\leq \frac{1}{\int_{\Omega} w d\mu} \sum_{i=1}^n \int_{\Omega} w_i \left| f - \frac{m+M}{2} \right| d\mu \\
&= \frac{1}{\int_{\Omega} w d\mu} \int_{\Omega} w \left| f - \frac{m+M}{2} \right| d\mu
\end{aligned}$$

and the last part of (2.17) is proved.  $\square$

If we use Lemma 2.3 for the discrete measure, we can state the following result:

**Lemma 2.11.** *Let  $\Phi : [a, b] \rightarrow \mathbb{R}$  be a convex function on the interval of real numbers  $[a, b]$ ,  $z_i \in [a, b]$ ,  $p_i \geq 0$ ,  $i \in \{1, \dots, n\}$  and  $\sum_{i=1}^n p_i = 1$ . Then*

$$\begin{aligned} 0 &\leq \sum_{i=1}^n p_i \Phi(z_i) - \Phi\left(\sum_{i=1}^n p_i z_i\right) \\ &\leq \left[\frac{\Phi(a) + \Phi(b)}{2} - \Phi\left(\frac{a+b}{2}\right)\right] \left(1 + \frac{2}{b-a} \left|\sum_{i=1}^n p_i z_i - \frac{a+b}{2}\right|\right) \\ &\leq 2 \left[\frac{\Phi(a) + \Phi(b)}{2} - \Phi\left(\frac{a+b}{2}\right)\right]. \end{aligned} \quad (2.19)$$

Using this lemma we can state and prove the following result as well:

**Theorem 2.12.** *Let  $\Phi : [m, M] \rightarrow \mathbb{R}$  be a convex function,  $f : \Omega \rightarrow [m, M]$  a  $\mu$ -measurable function such that  $f, \Phi \circ f \in L_w(\Omega, \mu)$ . Then for any  $\bar{w} \in \mathfrak{P}_n(w)$ ,  $n \geq 2$  we have*

$$\begin{aligned} 0 &\leq \psi(\Phi, f, \bar{w}) - \Phi\left(\frac{\int_{\Omega} f w d\mu}{\int_{\Omega} w d\mu}\right) \\ &\leq \left[\frac{\Phi(l(\bar{w})) + \Phi(L(\bar{w}))}{2} - \Phi\left(\frac{l(\bar{w}) + L(\bar{w})}{2}\right)\right] \\ &\quad \times \left(1 + \frac{2}{L(\bar{w}) - l(\bar{w})} \left|\frac{1}{\int_{\Omega} w d\mu} \int_{\Omega} w f d\mu - \frac{l(\bar{w}) + L(\bar{w})}{2}\right|\right) \\ &\leq 2 \left[\frac{\Phi(l(\bar{w})) + \Phi(L(\bar{w}))}{2} - \Phi\left(\frac{l(\bar{w}) + L(\bar{w})}{2}\right)\right], \end{aligned} \quad (2.20)$$

where  $l(\bar{w}), L(\bar{w})$  are defined by (2.13).

*Proof.* If we write the first two inequalities in (2.11) for

$$p_i = \frac{\int_{\Omega} w_i d\mu}{\int_{\Omega} w d\mu} > 0 \quad z_i = \frac{\int_{\Omega} f w_i d\mu}{\int_{\Omega} w_i d\mu}, \quad i \in \{1, \dots, n\}$$

and for  $a = l(\bar{w})$ ,  $b = L(\bar{w})$  as above we have

$$\begin{aligned}
0 &\leq \frac{1}{\int_{\Omega} w d\mu} \sum_{i=1}^n \left( \int_{\Omega} w_i d\mu \right) \Phi \left( \frac{\int_{\Omega} f w_i d\mu}{\int_{\Omega} w_i d\mu} \right) \\
&\quad - \Phi \left( \frac{\sum_{i=1}^n \frac{\int_{\Omega} w_i d\mu}{\int_{\Omega} w d\mu} \cdot \frac{\int_{\Omega} f w_i d\mu}{\int_{\Omega} w_i d\mu}}{\int_{\Omega} w d\mu} \right) \\
&\leq \left[ \frac{\Phi(l(\bar{w})) + \Phi(L(\bar{w}))}{2} - \Phi \left( \frac{l(\bar{w}) + L(\bar{w})}{2} \right) \right] \\
&\quad \times \left( 1 + \frac{2}{L(\bar{w}) - l(\bar{w})} \left| \sum_{i=1}^n \frac{\int_{\Omega} w_i d\mu}{\int_{\Omega} w d\mu} \frac{\int_{\Omega} f w_i d\mu}{\int_{\Omega} w_i d\mu} - \frac{l(\bar{w}) + L(\bar{w})}{2} \right| \right) \\
&= \left[ \frac{\Phi(l(\bar{w})) + \Phi(L(\bar{w}))}{2} - \Phi \left( \frac{l(\bar{w}) + L(\bar{w})}{2} \right) \right] \\
&\quad \times \left( 1 + \frac{2}{L(\bar{w}) - l(\bar{w})} \left| \frac{1}{\int_{\Omega} w d\mu} \int_{\Omega} \left( \sum_{i=1}^n w_i \right) f d\mu - \frac{l(\bar{w}) + L(\bar{w})}{2} \right| \right)
\end{aligned}$$

that proves the required inequalities in (2.20).  $\square$

### 3 Discrete Case and Some Applications

Let  $\Phi : [a, b] \rightarrow \mathbb{R}$  be a convex function on the interval of real numbers  $[a, b]$ ,  $x_k \in [a, b]$ ,  $w_k > 0$ ,  $k \in \{1, \dots, m\}$ . Let  $w_{ki} > 0$  for  $k \in \{1, \dots, m\}$ ,  $i \in \{1, \dots, n\}$  with  $m, n \geq 2$  and  $\sum_{i=1}^n w_{ki} = w_k$  for any  $k \in \{1, \dots, m\}$ .

We consider the functional associated with the matrix  $W := \{w_{ki}\}_{k \in \{1, \dots, m\}, i \in \{1, \dots, n\}}$

$$\psi(\Phi, \bar{x}, W) := \frac{1}{\sum_{k=1}^m w_k} \sum_{i=1}^n \Phi \left( \frac{\sum_{k=1}^m x_k w_{ki}}{\sum_{k=1}^m w_{ki}} \right) \sum_{k=1}^m w_{ki}, \quad (3.1)$$

where  $\bar{x} = (x_1, \dots, x_m) \in [a, b]^m$ .

Using the results from the previous section we have

$$\frac{\sum_{k=1}^m \Phi(x_k) w_k}{\sum_{k=1}^m w_k} \geq \psi(\Phi, \bar{x}, W) \geq \Phi \left( \frac{\sum_{k=1}^m x_k w_k}{\sum_{k=1}^m w_k} \right), \quad (3.2)$$

$$\begin{aligned}
0 &\leq \frac{\sum_{k=1}^m \Phi(x_k) w_k}{\sum_{k=1}^m w_k} - \psi(\Phi, \bar{x}, W) \\
&\leq \frac{\Phi'_-(b) - \Phi'_+(a)}{b-a} \\
&\times \frac{1}{\sum_{k=1}^m w_k} \sum_{i=1}^n \sum_{k=1}^m w_{ki} \left( b - \frac{\sum_{k=1}^m x_k w_{ki}}{\sum_{k=1}^m w_{ki}} \right) \left( \frac{\sum_{k=1}^m x_k w_{ki}}{\sum_{k=1}^m w_{ki}} - a \right) \\
&\leq \frac{\Phi'_-(b) - \Phi'_+(a)}{b-a} \left( b - \frac{\sum_{k=1}^m x_k w_k}{\sum_{k=1}^m w_k} \right) \left( \frac{\sum_{k=1}^m x_k w_k}{\sum_{k=1}^m w_k} - a \right),
\end{aligned} \tag{3.3}$$

and

$$\begin{aligned}
0 &\leq \frac{\sum_{k=1}^m \Phi(x_k) w_k}{\sum_{k=1}^m w_k} - \psi(\Phi, \bar{x}, W) \\
&\leq \left[ \frac{\Phi(a) + \Phi(b)}{2} - \Phi\left(\frac{a+b}{2}\right) \right] \\
&\times \left( 1 + \frac{2}{b-a} \sum_{i=1}^n \left| \frac{1}{\sum_{k=1}^m w_k} \sum_{k=1}^m w_{ki} \left( x_k - \frac{a+b}{2} \right) \right| \right) \\
&\leq \left[ \frac{\Phi(a) + \Phi(b)}{2} - \Phi\left(\frac{a+b}{2}\right) \right] \\
&\times \left( 1 + \frac{2}{b-a} \frac{1}{\sum_{k=1}^m w_k} \sum_{k=1}^m w_k \left| x_k - \frac{a+b}{2} \right| \right).
\end{aligned} \tag{3.4}$$

Define

$$l(W) := \min_{i \in \{1, \dots, n\}} \left\{ \frac{\sum_{k=1}^m w_{ki} x_k}{\sum_{k=1}^m w_{ki}} \right\}, \quad L(W) := \max_{i \in \{1, \dots, n\}} \left\{ \frac{\sum_{k=1}^m w_{ki} x_k}{\sum_{k=1}^m w_{ki}} \right\}. \tag{3.5}$$

Then we also have

$$\begin{aligned}
0 &\leq \psi(\Phi, \bar{x}, W) - \Phi\left(\frac{\sum_{k=1}^m x_k w_k}{\sum_{k=1}^m w_k}\right) \\
&\leq \frac{\Phi'_-(L(W)) - \Phi'_+(l(W))}{L(W) - l(W)} \\
&\times \left( L(W) - \frac{\sum_{k=1}^m x_k w_k}{\sum_{k=1}^m w_k} \right) \left( \frac{\sum_{k=1}^m x_k w_k}{\sum_{k=1}^m w_k} - l(W) \right) \\
&\leq \frac{1}{4} (L(W) - l(W)) [\Phi'_-(L(W)) - \Phi'_+(l(W))]
\end{aligned} \tag{3.6}$$

and

$$\begin{aligned}
0 &\leq \psi(\Phi, \bar{x}, W) - \Phi\left(\frac{\sum_{k=1}^m x_k w_k}{\sum_{k=1}^m w_k}\right) \\
&\leq \left[ \frac{\Phi(l(W)) + \Phi(L(W))}{2} - \Phi\left(\frac{l(W) + L(W)}{2}\right) \right] \\
&\quad \times \left( 1 + \frac{2}{L(W) - l(W)} \left| \frac{\sum_{k=1}^m x_k w_k}{\sum_{k=1}^m w_k} - \frac{l(W) + L(W)}{2} \right| \right) \\
&\leq 2 \left[ \frac{\Phi(l(W)) + \Phi(L(W))}{2} - \Phi\left(\frac{l(W) + L(W)}{2}\right) \right].
\end{aligned} \tag{3.7}$$

We consider the convex function  $\Phi : (0, \infty) \rightarrow (0, \infty)$ ,  $\Phi(t) = t^p$  with  $p \in (-\infty, 0) \cup (1, \infty)$ . Then

$$\begin{aligned}
\psi(\Phi, \bar{x}, W) &:= \frac{1}{\sum_{k=1}^m w_k} \sum_{i=1}^n \left( \frac{\sum_{k=1}^m x_k w_{ki}}{\sum_{k=1}^m w_{ki}} \right)^p \sum_{k=1}^m w_{ki} \\
&= \frac{1}{\sum_{k=1}^m w_k} \sum_{i=1}^n \left( \sum_{k=1}^m x_k w_{ki} \right)^p \left( \sum_{k=1}^m w_{ki} \right)^{1-p}
\end{aligned} \tag{3.8}$$

where  $x_k > 0$ ,  $w_k > 0$ ,  $w_{ki} > 0$  for  $k \in \{1, \dots, m\}$ ,  $i \in \{1, \dots, n\}$  with  $m, n \geq 2$  and  $\sum_{i=1}^n w_{ki} = w_k$  for any  $k \in \{1, \dots, m\}$ .

From (3.2) we have

$$\begin{aligned}
\frac{\sum_{k=1}^m x_k^p w_k}{\sum_{k=1}^m w_k} &\geq \frac{1}{\sum_{k=1}^m w_k} \sum_{i=1}^n \left( \sum_{k=1}^m x_k w_{ki} \right)^p \left( \sum_{k=1}^m w_{ki} \right)^{1-p} \\
&\geq \left( \frac{\sum_{k=1}^m x_k w_k}{\sum_{k=1}^m w_k} \right)^p.
\end{aligned} \tag{3.9}$$

If we set  $a = \min \{x_k\}_{k \in \{1, \dots, m\}}$  and  $b = \max \{x_k\}_{k \in \{1, \dots, m\}}$  then from (3.3) and (3.4) we have

$$\begin{aligned}
0 &\leq \frac{\sum_{k=1}^m x_k^p w_k}{\sum_{k=1}^m w_k} - \frac{1}{\sum_{k=1}^m w_k} \sum_{i=1}^n \left( \sum_{k=1}^m x_k w_{ki} \right)^p \left( \sum_{k=1}^m w_{ki} \right)^{1-p} \\
&\leq p \frac{b^{p-1} - a^{p-1}}{b - a} \\
&\quad \times \frac{1}{\sum_{k=1}^m w_k} \sum_{i=1}^n \sum_{k=1}^m w_{ki} \left( b - \frac{\sum_{k=1}^m x_k w_{ki}}{\sum_{k=1}^m w_{ki}} \right) \left( \frac{\sum_{k=1}^m x_k w_{ki}}{\sum_{k=1}^m w_{ki}} - a \right) \\
&\leq p \frac{b^{p-1} - a^{p-1}}{b - a} \left( b - \frac{\sum_{k=1}^m x_k w_k}{\sum_{k=1}^m w_k} \right) \left( \frac{\sum_{k=1}^m x_k w_k}{\sum_{k=1}^m w_k} - a \right),
\end{aligned} \tag{3.10}$$

and

$$\begin{aligned}
0 &\leq \frac{\sum_{k=1}^m x_k^p w_k}{\sum_{k=1}^m w_k} - \frac{1}{\sum_{k=1}^m w_k} \sum_{i=1}^n \left( \sum_{k=1}^m x_k w_{ki} \right)^p \left( \sum_{k=1}^m w_{ki} \right)^{1-p} \\
&\leq \left[ \frac{a^p + b^p}{2} - \left( \frac{a+b}{2} \right)^p \right] \\
&\times \left( 1 + \frac{2}{b-a} \sum_{i=1}^n \left| \frac{1}{\sum_{k=1}^m w_k} \sum_{k=1}^m w_{ki} \left( x_k - \frac{a+b}{2} \right) \right| \right) \\
&\leq \left[ \frac{a^p + b^p}{2} - \left( \frac{a+b}{2} \right)^p \right] \\
&\times \left( 1 + \frac{2}{b-a} \frac{1}{\sum_{k=1}^m w_k} \sum_{k=1}^m w_k \left| x_k - \frac{a+b}{2} \right| \right).
\end{aligned} \tag{3.11}$$

If  $l(W)$  and  $L(W)$  are defined as in (3.5), then from (3.6) and (3.7) we also have

$$\begin{aligned}
0 &\leq \frac{1}{\sum_{k=1}^m w_k} \sum_{i=1}^n \left( \sum_{k=1}^m x_k w_{ki} \right)^p \left( \sum_{k=1}^m w_{ki} \right)^{1-p} - \left( \frac{\sum_{k=1}^m x_k w_k}{\sum_{k=1}^m w_k} \right)^p \\
&\leq p \frac{(L(W))^{p-1} - (l(W))^{p-1}}{L(W) - l(W)} \\
&\times \left( L(W) - \frac{\sum_{k=1}^m x_k w_k}{\sum_{k=1}^m w_k} \right) \left( \frac{\sum_{k=1}^m x_k w_k}{\sum_{k=1}^m w_k} - l(W) \right) \\
&\leq \frac{1}{4} (L(W) - l(W)) \left[ (L(W))^{p-1} - (l(W))^{p-1} \right]
\end{aligned} \tag{3.12}$$

and

$$\begin{aligned}
0 &\leq \frac{1}{\sum_{k=1}^m w_k} \sum_{i=1}^n \left( \sum_{k=1}^m x_k w_{ki} \right)^p \left( \sum_{k=1}^m w_{ki} \right)^{1-p} - \left( \frac{\sum_{k=1}^m x_k w_k}{\sum_{k=1}^m w_k} \right)^p \\
&\leq \left[ \frac{(l(W))^p + (L(W))^p}{2} - \left( \frac{l(W) + L(W)}{2} \right)^p \right] \\
&\times \left( 1 + \frac{2}{L(W) - l(W)} \left| \frac{\sum_{k=1}^m x_k w_k}{\sum_{k=1}^m w_k} - \frac{l(W) + L(W)}{2} \right| \right) \\
&\leq 2 \left[ \frac{(l(W))^p + (L(W))^p}{2} - \left( \frac{l(W) + L(W)}{2} \right)^p \right].
\end{aligned} \tag{3.13}$$

Further on, consider the convex function  $\Phi : (0, \infty) \rightarrow (0, \infty)$ ,  $\Phi(t) = -\ln t$  then from (3.2) for  $x_k > 0$ ,  $w_k > 0$ ,  $w_{ki} > 0$  for  $k \in \{1, \dots, m\}$ ,  $i \in \{1, \dots, n\}$  with  $m$ ,

$n \geq 2$  and  $\sum_{i=1}^n w_{ki} = w_k$  for any  $k \in \{1, \dots, m\}$  we get

$$\begin{aligned} \left( \prod_{k=1}^m x_k^{w_k} \right)^{\frac{1}{\sum_{k=1}^m w_k}} &\leq \left( \prod_{i=1}^n \left( \frac{\sum_{k=1}^m x_k w_{ki}}{\sum_{k=1}^m w_{ki}} \right)^{\sum_{k=1}^m w_{ki}} \right)^{\frac{1}{\sum_{k=1}^m w_k}} \\ &\leq \frac{\sum_{k=1}^m x_k w_k}{\sum_{k=1}^m w_k}. \end{aligned} \quad (3.14)$$

By (3.3) and (3.4) we have

$$\begin{aligned} 1 &\leq \frac{\left( \prod_{i=1}^n \left( \frac{\sum_{k=1}^m x_k w_{ki}}{\sum_{k=1}^m w_{ki}} \right)^{\sum_{k=1}^m w_{ki}} \right)^{\frac{1}{\sum_{k=1}^m w_k}}}{\left( \prod_{k=1}^m x_k^{w_k} \right)^{\frac{1}{\sum_{k=1}^m w_k}}} \\ &\leq \exp \left[ \frac{1}{ba \sum_{k=1}^m w_k} \sum_{i=1}^n \sum_{k=1}^m w_{ki} \left( b - \frac{\sum_{k=1}^m x_k w_{ki}}{\sum_{k=1}^m w_{ki}} \right) \left( \frac{\sum_{k=1}^m x_k w_{ki}}{\sum_{k=1}^m w_{ki}} - a \right) \right] \\ &\leq \exp \left[ \frac{1}{ab} \left( b - \frac{\sum_{k=1}^m x_k w_k}{\sum_{k=1}^m w_k} \right) \left( \frac{\sum_{k=1}^m x_k w_k}{\sum_{k=1}^m w_k} - a \right) \right] \end{aligned} \quad (3.15)$$

and

$$\begin{aligned} 1 &\leq \frac{\left( \prod_{i=1}^n \left( \frac{\sum_{k=1}^m x_k w_{ki}}{\sum_{k=1}^m w_{ki}} \right)^{\sum_{k=1}^m w_{ki}} \right)^{\frac{1}{\sum_{k=1}^m w_k}}}{\left( \prod_{k=1}^m x_k^{w_k} \right)^{\frac{1}{\sum_{k=1}^m w_k}}} \\ &\leq \left( \frac{a+b}{2\sqrt{ab}} \right)^{1 + \frac{2}{b-a} \sum_{i=1}^n \left| \frac{1}{\sum_{k=1}^m w_{ki}} \sum_{k=1}^m w_{ki} (x_k - \frac{a+b}{2}) \right|} \\ &\leq \left( \frac{a+b}{2\sqrt{ab}} \right)^{1 + \frac{2}{b-a} \frac{1}{\sum_{k=1}^m w_k} \sum_{k=1}^m w_k |x_k - \frac{a+b}{2}|}. \end{aligned} \quad (3.16)$$

From (3.6) and (3.7) we also have

$$\begin{aligned} 1 &\leq \frac{\frac{1}{\sum_{k=1}^m w_k} \sum_{k=1}^m x_k w_k}{\left( \prod_{i=1}^n \left( \frac{\sum_{k=1}^m x_k w_{ki}}{\sum_{k=1}^m w_{ki}} \right)^{\sum_{k=1}^m w_{ki}} \right)^{\frac{1}{\sum_{k=1}^m w_k}}} \\ &\leq \exp \left[ \frac{1}{L(W)l(W)} \left( L(W) - \frac{\sum_{k=1}^m x_k w_k}{\sum_{k=1}^m w_k} \right) \left( \frac{\sum_{k=1}^m x_k w_k}{\sum_{k=1}^m w_k} - l(W) \right) \right] \\ &\leq \frac{1}{4} \frac{(L(W) - l(W))^2}{L(W)l(W)} \end{aligned} \quad (3.17)$$

and

$$\begin{aligned}
1 &\leq \frac{\frac{1}{\sum_{k=1}^m w_k} \sum_{k=1}^m x_k w_k}{\left( \prod_{i=1}^n \left( \frac{\sum_{k=1}^m x_k w_{ki}}{\sum_{k=1}^m w_{ki}} \right)^{\sum_{k=1}^m w_{ki}} \right)^{\frac{1}{\sum_{k=1}^m w_k}}} & (3.18) \\
&\leq \left( \frac{l(W) + L(W)}{2\sqrt{l(W)L(W)}} \right)^{1 + \frac{2}{L(W)-l(W)} \left| \frac{\sum_{k=1}^m x_k w_k}{\sum_{k=1}^m w_k} - \frac{l(W)+L(W)}{2} \right|} \\
&\leq \left( \frac{l(W) + L(W)}{2\sqrt{l(W)L(W)}} \right)^2.
\end{aligned}$$

## 4 Applications for Univariate Functions

Let  $\Phi : [m, M] \rightarrow \mathbb{R}$  be a convex function and  $f : [0, \pi] \rightarrow [m, M]$  an integrable function. Since  $\sin^2 t + \cos^2 t = 1$  for any  $t \in [0, \pi]$  then  $\bar{w} = (\sin^2, \cos^2)$  is a partition of the unity. We then have

$$\begin{aligned}
&\psi(\Phi, f, \bar{w}) \\
&:= \frac{1}{\pi} \left[ \Phi \left( \frac{\int_0^\pi f(t) \sin^2 t dt}{\int_0^\pi \sin^2 t dt} \right) \int_0^\pi \sin^2 t dt + \Phi \left( \frac{\int_0^\pi f(t) \cos^2 t dt}{\int_0^\pi \cos^2 t dt} \right) \int_0^\pi \cos^2 t dt \right] \\
&= \frac{1}{2} \left[ \Phi \left( \frac{2}{\pi} \int_0^\pi f(t) \sin^2 t dt \right) + \Phi \left( \frac{2}{\pi} \int_0^\pi f(t) \cos^2 t dt \right) \right].
\end{aligned}$$

By the inequality (2.1) we have

$$\begin{aligned}
\frac{\int_0^\pi (\Phi \circ f)(t) dt}{\pi} &\geq \frac{1}{2} \left[ \Phi \left( \frac{2}{\pi} \int_0^\pi f(t) \sin^2 t dt \right) + \Phi \left( \frac{2}{\pi} \int_0^\pi f(t) \cos^2 t dt \right) \right] & (4.1) \\
&\geq \Phi \left( \frac{\int_0^\pi f(t) dt}{\pi} \right),
\end{aligned}$$

while from (2.10) we have

$$\begin{aligned}
0 &\leq \frac{\int_0^\pi (\Phi \circ f)(t) dt}{\pi} & (4.2) \\
&\quad - \frac{1}{2} \left[ \Phi \left( \frac{2}{\pi} \int_0^\pi f(t) \sin^2 t dt \right) + \Phi \left( \frac{2}{\pi} \int_0^\pi f(t) \cos^2 t dt \right) \right]
\end{aligned}$$

$$\begin{aligned}
&\leq \frac{1}{2} \frac{\Phi'_-(M) - \Phi'_+(m)}{M - m} \\
&\left[ \left( M - \frac{2}{\pi} \int_0^\pi f(t) \sin^2 t dt \right) \left( \frac{2}{\pi} \int_0^\pi f(t) \sin^2 t dt - m \right) \right. \\
&+ \left. \left( M - \frac{2}{\pi} \int_0^\pi f(t) \cos^2 t dt \right) \left( \frac{2}{\pi} \int_0^\pi f(t) \cos^2 t dt - m \right) \right] \\
&\leq \frac{\Phi'_-(M) - \Phi'_+(m)}{M - m} \left( M - \frac{\int_0^\pi f(t) dt}{\pi} \right) \left( \frac{\int_0^\pi f(t) dt}{\pi} - m \right).
\end{aligned}$$

Now, let

$$\begin{aligned}
l(\bar{w}) &: = \frac{2}{\pi} \min \left\{ \int_0^\pi f(t) \sin^2 t dt, \int_0^\pi f(t) \cos^2 t dt \right\} \\
&= \frac{1}{\pi} \left[ \int_0^\pi f(t) dt - \left| \int_0^\pi f(t) \cos 2t dt \right| \right]
\end{aligned}$$

and

$$\begin{aligned}
L(\bar{w}) &: = \frac{2}{\pi} \max \left\{ \int_0^\pi f(t) \sin^2 t dt, \int_0^\pi f(t) \cos^2 t dt \right\} \\
&= \frac{1}{\pi} \left[ \int_0^\pi f(t) dt + \left| \int_0^\pi f(t) \cos 2t dt \right| \right]
\end{aligned}$$

then by (2.15) we have

$$\begin{aligned}
0 &\leq \frac{1}{2} \left[ \Phi \left( \frac{2}{\pi} \int_0^\pi f(t) \sin^2 t dt \right) + \Phi \left( \frac{2}{\pi} \int_0^\pi f(t) \cos^2 t dt \right) \right] \\
&- \Phi \left( \frac{\int_0^\pi f(t) dt}{\pi} \right) \\
&\leq \frac{1}{2\pi} \left| \int_0^\pi f(t) \cos 2t dt \right| \\
&\times \left[ \Phi'_- \left( \frac{1}{\pi} \left[ \int_0^\pi f(t) dt + \left| \int_0^\pi f(t) \cos 2t dt \right| \right] \right) \right. \\
&- \left. \Phi'_+ \left( \frac{1}{\pi} \left[ \int_0^\pi f(t) dt - \left| \int_0^\pi f(t) \cos 2t dt \right| \right] \right) \right].
\end{aligned} \tag{4.3}$$

Similar inequalities may be obtained if someone would also use the inequalities (2.17) and (2.20). The details are omitted.

## References

- [1] S. Abramovich, S. Ivelić and J. Pečarić, Generalizations of JensenSteffensen and related integral inequalities for superquadratic functions. *Cent. Eur. J. Math.* **8** (2010), no. 5, 937–949.
- [2] S. S. Dragomir, An improvement of Jensen's inequality, *Bull. Math. Soc. Sci. Math. Roumanie*, **34 (82)** (1990), No. 4, 291-296.
- [3] S. S. Dragomir, Some refinements of Ky Fan's inequality, *J. Math. Anal. Appl.*, **163** (2) (1992), 317-321.
- [4] S. S. Dragomir, Some refinements of Jensen's inequality, *J. Math. Anal. Appl.*, **168** (2) (1992), 518-522.
- [5] S. S. Dragomir, A further improvement of Jensen's inequality, *Tamkang J. Math.*, **25** (1) (1994), 29-36.
- [6] S. S. Dragomir, A new improvement of Jensen's inequality, *Indian J. Pure and Appl. Math.*, **26** (10) (1995), 959-968.
- [7] S. S. Dragomir, A refinement of Jensen's inequality with applications for  $f$ -divergence measures, *Taiwanese J. Math.* **14** (2010), no. 1, 153–164. Preprint, *Res. Rep. Coll.* **10** (2007), Supp., Article 15.
- [8] S. S. Dragomir, A new refinement of Jensen's inequality in linear spaces with applications. *Math. Comput. Modelling* **52** (2010), no. 9-10, 1497–1505.
- [9] S. S. Dragomir, Some reverses of the Jensen inequality with applications. *Bull. Aust. Math. Soc.* **87** (2013), no. 2, 177–194.
- [10] S. S. Dragomir and N. M. Ionescu, Some converse of Jensen's inequality and applications. *Rev. Anal. Numér. Théor. Approx.* **23** (1994), no. 1, 71–78. MR1325895 (96c:26012).
- [11] S. S. Dragomir and C. J. Goh, A counterpart of Jensen's discrete inequality for differentiable convex mappings and applications in information theory, *Math. Comput. Modelling*, **24** (1996), No. 2, 1-11.
- [12] S. S. Dragomir, J. Pečarić and L. E. Persson, Properties of some functionals related to Jensen's inequality, *Acta Math. Hung.*, **70** (1-2) (1996), 129-143.
- [13] S. Khalid and J. Pečarić, On the refinements of the integral JensenSteffensen inequality. *J. Inequal. Appl.* **2013**, 2013:20, 18 pp.

- [14] J. Pečarić and S. S. Dragomir, A refinements of Jensen inequality and applications, *Studia Univ. Babeş-Bolyai, Mathematica*, **24** (1) (1989), 15-19.
- [15] F. Qi and M.-L. Yang, Comparisons of two integral inequalities with Hermite-Hadamard-Jensen's integral inequality. *Int. J. Appl. Math. Sci.* **3** (2006), no. 1, 83–88.
- [16] Z. Y. Song, Discussion on the integralttype Jensen inequality for P-convex functions. (Chinese) *Pure Appl. Math. (Xi'an)* **28** (2012), no. 1, 36–40.

S. S. Dragomir

<sup>1</sup>Mathematics, College of Engineering & Science  
Victoria University, PO Box 14428  
Melbourne City, MC 8001, Australia.

<sup>2</sup>DST-NRF Centre of Excellence in the  
Mathematical and Statistical Sciences  
School of Computer Science & Applied Mathematics,  
University of the Witwatersrand, Private Bag 3,  
Johannesburg 2050, South Africa  
E-mail: sever.dragomir@vu.edu.au