

p -Minkowski Inequalities for Mixed Intersection Bodies

Chang-jian ZHAO* and Mihály BENCZE

Abstract. In this paper, we establish p -Minkowski inequality for mixed intersection bodies, which is a general form of the Minkowski inequality for mixed intersection bodies.

Keywords: dual mixed volumes, mixed intersection bodies.

Mathematics Subject Classification (2000): Primary 52A40.

One might say the history of intersection bodies began with the paper [1] of Busemann. Intersection bodies were first explicitly defined and named by Lutwak [2]. It was here that the duality between intersection bodies and projection bodies was first made clear. Despite the considerable ingenuity of earlier attacks on the Busemann-Petty problem, it seems fair to say that the work [2] of Lutwak represents the beginning of its eventual solution. In [2], Lutwak also showed that if a convex body is sufficiently smooth and not an intersection body, then there exists a centred star body such that the conditions of Busemann-Petty problem holds, but the result inequality is reversed. Following Lutwak, the intersection body of order i of a star body is introduced by Zhang [3]. It follows from this definition that every intersection body of order i of a star body is an intersection body of a star body, and vice versa. As Zhang observes, the new definition of intersection body allows a more appealing formulation, namely: The Busemann-Petty problem has a positive answer in n -dimensional Euclidean space if and only if each centered convex body is an intersection body. The intersection body plays an essential role in Busemann's theory [4] of area in Minkowski spaces. The intersection body also is an important matter of the Brunn-Minkowski theory.

In recent years some author including Ball [5-6], Bourgain [7], Gardner [8-10], Schneider [11] and Lutwak [12-18] et al have given considerable attention to the Brunn-Minkowski theory and their various generalizations. The purpose of this paper first is to establish the Minkowski inequality for the dual Quermassintegral sum, which is a generalization of the Minkowski inequality for mixed intersection bodies. Then, the Brunn-Minkowski inequality and the Aleksandrov-Fenchel inequality for mixed intersection bodies are proved and some related results also be given. In this work we shall derive, for intersection bodies, all the analogous inequalities for Lutwak's mixed projection body inequalities [15]. Thus, this work may be seen as presenting additional evidence of the natural duality between intersection and projection bodies.

* Author to whom correspondence should be addressed. Research is supported by Zhejiang Provincial Natural Science Foundation of China(Y605065), Foundation of the Education Department of Zhejiang Province of China(20050392).

1 Notation and Preliminaries

The setting for this paper is n -dimensional Euclidean space \mathbb{R}^n ($n > 2$). Let \mathbb{C}^n denote the set of non-empty convex figures (compact, convex subsets) and \mathcal{K}^n denote the subset of \mathbb{C}^n consisting of all convex bodies (compact, convex subsets with non-empty interiors) in \mathbb{R}^n . We reserve the letter u for unit vectors, and the letter B is reserved for the unit ball centered at the origin. The surface of B is S^{n-1} . For $u \in S^{n-1}$, let E_u denote the hyperplane, through the origin, that is orthogonal to u . We will use K^u to denote the image of K under an orthogonal projection onto the hyperplane E_u . We use $V(K)$ for the n -dimensional volume of convex body K . The support function of $K \in \mathcal{K}^n$, $h(K, \cdot)$, defined on \mathbb{R}^n by $h(K, \cdot) = \text{Max}\{x \cdot y : y \in K\}$. Let δ denote the Hausdorff metric on \mathcal{K}^n ; i.e., for $K, L \in \mathcal{K}^n$, $\delta(K, L) = |h_K - h_L|_\infty$, where $|\cdot|_\infty$ denotes the sup-norm on the space of continuous functions, $C(S^{n-1})$.

Associated with a compact subset K of \mathbb{R}^n , which is star-shaped with respect to the origin, its radial function $\rho(K, \cdot) : S^{n-1} \rightarrow \mathbb{R}$, defined for $u \in S^{n-1}$, by $\rho(K, u) = \text{Max}\{\lambda \geq 0 : \lambda u \in K\}$. If $\rho(K, \cdot)$ is positive and continuous, K will be called a star body. Let φ^n denote the set of star bodies in \mathbb{R}^n .

1.1 Dual mixed volumes

If $K_1, \dots, K_r \in \varphi^n$ and $\lambda_1, \dots, \lambda_r \in \mathbb{R}$, then the radial Minkowski linear combination, $\lambda_1 K_1 \tilde{+} \dots \tilde{+} \lambda_r K_r$, is defined by $\lambda_1 K_1 \tilde{+} \dots \tilde{+} \lambda_r K_r = \{\lambda_1 x_1 \tilde{+} \dots \tilde{+} \lambda_r x_r : x_i \in K_i\}$.

The following property will be used later. If $K, L \in \varphi^n$ and $\lambda, \mu \geq 0$

$$\rho(\lambda K \tilde{+} \mu L, \cdot) = \lambda \rho(K, \cdot) + \mu \rho(L, \cdot) \quad (1.1.1)$$

For $K_1, \dots, K_r \in \varphi^n$ and $\lambda_1, \dots, \lambda_r \geq 0$, the volume of the radial Minkowski linear combination $\lambda_1 K_1 \tilde{+} \dots \tilde{+} \lambda_r K_r$ is a homogeneous n th-degree polynomial in the λ_i [11],

$$V(\lambda_1 K_1 \tilde{+} \dots \tilde{+} \lambda_r K_r) = \sum \tilde{V}_{i_1, \dots, i_n} \lambda_{i_1} \dots \lambda_{i_n} \quad (1.1.2)$$

where the sum is taken over all n -tuples (i_1, \dots, i_n) whose entries are positive integers not exceeding r . If we require the coefficients of the polynomial in (1.1.2) to be symmetric in their arguments, then they are uniquely determined. The coefficient $\tilde{V}_{i_1, \dots, i_n}$ is nonnegative and depends only on the bodies K_{i_1}, \dots, K_{i_n} . It is written as $\tilde{V}(K_{i_1}, \dots, K_{i_n})$ and is called the *dual mixed volume* of K_{i_1}, \dots, K_{i_n} . If $K_1 = \dots = K_{n-i} = K$, $K_{n-i+1} = \dots = K_n = L$, the dual mixed volume is written as $\tilde{V}_i(K, L)$. The dual mixed volume $\tilde{V}_i(K, B)$ is written as $\tilde{W}_i(K)$.

Dual Quermassintegral is special case of the p -th dual volume:

$$\tilde{V}_p(K) = \frac{1}{n} \int_{S^{n-1}} \rho(K, u)^p dS(u), \quad -\infty < p < \infty. \quad (1.1.3)$$

Taking for $p = n - i$ in $\tilde{V}_p(K)$, $\tilde{V}_p(K)$ changes to the well know dual Quermassintegral $\tilde{W}_i(K)$. Taking for $p = n$ in $\tilde{V}_p(K)$, $\tilde{V}_p(K)$ changes to the well know general volume $V_i(K)$.

If $K_i \in \varphi^n$ ($i = 1, 2, \dots, n-1$), then the dual mixed volume of $K_i \cap E_u$ ($i = 1, 2, \dots, n-1$) will be denoted by $\tilde{v}(K_1 \cap E_u, \dots, K_{n-1} \cap E_u)$. If $K_1 = \dots = K_{n-1-i} = K$ and $K_{n-i} =$

$\dots = K_{n-1} = L$, then $\tilde{v}(K_1 \cap E_u, \dots, K_{n-1} \cap E_u)$ is written $\tilde{v}_i(K \cap E_u, L \cap E_u)$. If $L = B$, then $\tilde{v}_i(K \cap E_u, B \cap E_u)$ is written $\tilde{w}_i(K \cap E_u)$.

1.2 Intersection bodies

For $K \in \varphi^n$, there is a unique star body IK whose radial function satisfies for $u \in S^{n-1}$,

$$\rho(IK, u) = v(K \cap E_u), \tag{1.2.1}$$

It is called the *intersection bodies* of K . From a result of Busemann, it follows that IK is a convex if K is convex and centrally symmetric with respect to the origin. Clearly any intersection body is centred.

Volume of the intersection bodies is given by $V(IK) = \frac{1}{n} \int_{S^{n-1}} v(K \cap E_u)^n dS(u)$.

The mixed intersection bodies of $K_1, \dots, K_{n-1} \in \varphi^n$, $I(K_1, \dots, K_{n-1})$, whose radial function is defined by

$$\rho(I(K_1, \dots, K_{n-1}), u) = \tilde{v}(K_1 \cap E_u, \dots, K_{n-1} \cap E_u), \tag{1.2.2}$$

where \tilde{v} is $(n - 1)$ -dimensional dual mixed volume.

If $K \in \varphi^n$ with $\rho(K, u) \in C(S^{n-1})$, and $i \in \mathbb{R}$ is positive, the *intersection body of order i* of K is the centered star body $I_i K$ such that^[3] $\rho(I_i K) = \frac{1}{n-1} \int_{S^{n-1}} \rho(K, u)^{n-i-1} dS(u)$, for $u \in S^{n-1}$, where $I_i K = I(\underbrace{K, \dots, K}_{n-i-1}, \underbrace{B, \dots, B}_i)$.

If $K_1 = \dots = K_{n-i-1} = K, K_{n-i} = \dots = K_{n-1} = L$, then $I(K_1, \dots, K_{n-1})$ is written as $I_i(K, L)$. If $L = B$, then $I_i(K, L)$ is written as $I_i K$ is called the i th intersection body of K . For $I_0 K$ simply write IK . The term is introduced by Zhang [3].

2 Main results

2.0 Lemmas

Lemma A *If $K, L \in \varphi^n$, $-\infty < p < \infty$ and $0 < j < n - 1$ then*

$$\begin{aligned} \tilde{V}_p(IK) &= \frac{1}{n} \int_{S^{n-1}} v(K \cap E_u)^p dS(u); \\ \tilde{V}_p(I_j K) &= \frac{1}{n} \int_{S^{n-1}} \tilde{w}_j(K \cap E_u)^p dS(u); \\ \tilde{V}_p(I_j(K, L)) &= \frac{1}{n} \int_{S^{n-1}} \tilde{v}_j(K \cap E_u, L \cap E_u)^p dS(u). \end{aligned}$$

To prove this we use (1.1.3) in conjunction with the fact (1.2.2).

Lemma B^[14] *If $K_1, \dots, K_n \in \varphi^n$, then*

$$\tilde{V}(K_1, \dots, K_n)^r \leq \prod_{j=1}^r \tilde{V}(\underbrace{K_j, \dots, K_j}_r, K_{r+1}, \dots, K_n).$$

with equality if and only if K_1, \dots, K_n are all dilations of each other.

We shall need the following trivial elementary inequality:

Lemma C *If $a, b \geq 0$ and $c, d > 0$ then for $0 < p < 1$*

$$(a + b)^p (c + d)^{1-p} \geq a^p c^{p-1} + b^p d^{p-1},$$

with equality if and only if $ad = bc$.

2.1 The Minkowski inequality for dual Quermassintegral sum of mixed intersection bodies

In [19], Leng introduce the concept of *i-Quermassintegral difference function* of convex bodies as follows. If $K, D \in \mathcal{K}^n$ and $D \subset K$, then *i-Quermassintegral difference function* of convex bodies K and D , $D_{w_i}(K, D)$, was defined by

$$D_{w_i}(K, D) = W_i(K) - W_i(D) \quad (0 \leq i \leq n-1).$$

In the section, we first introduce a new concept, *dual Quermassintegral sum function*, as follows.

If $K, D \in \varphi^n$, then the p -dual volume sum function of star bodies K and D , $S_{\tilde{v}_p}(K, D)$, is defined by

$$S_{\tilde{v}_p}(K, D) = \tilde{V}_p(K) + \tilde{V}_p(D).$$

When $p = n$, we have $S_v(K, D) = V(K) + V(D)$, which is called the *dual volume sum function* of star bodies K and L .

The following Minkowski inequality for mixed intersection bodies will be established: If $K, L \in \varphi^n$, and $0 \leq i < n$ and $0 < j < n-1$, then

$$\tilde{W}_i(I_j(K, L))^{n-1} \leq \tilde{W}_i(IK)^{n-j-1} \tilde{W}_i(IL)^j, \quad (2.1.0)$$

with equality if and only if K and L are dilates.

This is just the special case $D = D' = \emptyset$ of:

Theorem 2.1.1 *If $K, L, D, D' \in \varphi^n$. Let D' is a dilates copy of D , and $-\infty < p < \infty$ and $0 < j < n-1$, then*

$$S_{\tilde{v}_p}(I_j(K, L), I_j(D, D'))^{n-1} \leq S_{\tilde{v}_p}(IK, ID)^{n-j-1} S_{\tilde{v}_p}(IL, ID')^j, \quad (2.1.1)$$

with equality if and only if K and L are dilates.

Proof In view of the special case of Lemma B, we obtain that

$$\tilde{v}_j(K \cap E_u, L \cap E_u) \leq v(K \cap E_u)^{\frac{(n-j-1)}{n-1}} v(L \cap E_u)^{\frac{j}{n-1}}. \quad (2.1.2)$$

with equality if and only if $K \cap E_u$ and $L \cap E_u$ are dilates, it follows if and only if K and L are dilates^[20].

From Lemma A, (2.1.2) and in view of Minkowski inequality for integral^[21], we have $j < n-1$

$$n\tilde{V}_p(I_j(K, L)) = (\|\tilde{v}_j(K \cap E_u, L \cap E_u)\|_p)^p$$

$$\begin{aligned}
 &\leq \left(\|v(K \cap E_u)^{\frac{n-j-1}{n-1}} v(L \cap E_u)^{\frac{j}{n-1}}\|_p \right)^p \\
 &\leq (\|v(K \cap E_u)\|_p)^{\frac{p(n-j-1)}{n-1}} (\|v(K \cap E_u)\|_p)^{\frac{jp}{n-1}} \\
 &= (n\tilde{V}_p(IK))^{\frac{(n-j-1)}{n-1}} (n\tilde{V}_p(IL))^{\frac{j}{n-1}} \\
 &= n\tilde{V}_p(IK)^{\frac{(n-j-1)}{n-1}} \tilde{V}_p(IL)^{\frac{j}{n-1}}. \tag{2.1.3}
 \end{aligned}$$

In view of the conditions of (2.1.2) and Minkowski inequality for integral, it follows that the equality holds if and only if K and L are dilates.

Moreover, we consider the case of $j = n - 1$ of the inequality (2.1.3). If $j = n - 1$, inequality (2.1.3) reduces to

$$\tilde{W}_{n-1}(I_j(K, L))^{n-1} \leq \tilde{W}_{n-1}(IK)^{n-j-1} \tilde{W}_{n-1}(IL)^j. \tag{*}$$

From Lemma A, (*) changes to

$$\begin{aligned}
 &\left(\int_{S^{n-1}} \tilde{v}_j(K \cap E_u, L \cap E_u) dS(u) \right)^{n-1} \\
 &\leq \left(\int_{S^{n-1}} v(K \cap E_u) dS(u) \right)^{n-j-1} \left(\int_{S^{n-1}} v(L \cap E_u) dS(u) \right)^j. \tag{**}
 \end{aligned}$$

On the other hand, integrating two sides of (2.1.2) and in view of Hölder inequality for integral, we obtain that

$$\begin{aligned}
 \int_{S^{n-1}} \tilde{v}_j(K \cap E_u, L \cap E_u) dS(u) &\leq \int_{S^{n-1}} v(K \cap E_u)^{\frac{n-j-1}{n-1}} v(L \cap E_u)^{\frac{j}{n-1}} dS(u) \\
 &\leq \left(\int_{S^{n-1}} v(K \cap E_u) dS(u) \right)^{\frac{n-j-1}{n-1}} \left(\int_{S^{n-1}} v(L \cap E_u) dS(u) \right)^{\frac{j}{n-1}}.
 \end{aligned}$$

Moreover, from the inequality (2.1.3), we obtain that

$$\tilde{V}_p(I_j(K, L))^{n-1} \leq \tilde{V}_p(IK)^{n-j-1} \tilde{V}_p(IL)^j,$$

with equality if and only if K and L are dilates, and

$$\tilde{V}_p(I_j(D, D'))^{n-1} = \tilde{V}_p(ID)^{n-j-1} \tilde{V}_p(ID')^j.$$

Hence, from the inequality in Lemma C, we have

$$\begin{aligned}
 &S_{\tilde{v}_p}(I_j(K, L), I_j(D, D')) \\
 &\leq \tilde{V}_p(IK)^{(n-j-1)/(n-1)} \tilde{V}_p(IL)^{j/(n-1)} + \tilde{V}_p(ID)^{(n-j-1)/(n-1)} \tilde{V}_p(ID')^{j/(n-1)} \\
 &\leq S_{\tilde{v}_p}(IK, ID)^{n-j-1} S_{\tilde{v}_p}(IL, ID')^j.
 \end{aligned}$$

The proof of Theorem 2.1.1 is complete.

Remark 2.1.1 Let D and D' be single point and take $p = n - i$ to (2.1.1), (2.1.1) changes to inequality (2.1.0)

$$\tilde{W}_i(I_1(K, L))^{n-1} \leq \tilde{W}_i(IK)^{n-2} \tilde{W}_i(IL),$$

with equality if and only if K and L are dilates.

This is just a dual form of the following inequality which was given by Lutwak [15].

The Minkowski inequality for mixed projection bodies If $K, L \in \mathcal{K}^n$, and $0 \leq i < n$, then

$$W_i(\Pi_1(K, L))^{n-1} \geq W_i(\Pi K)^{n-2} W_i(\Pi L),$$

with equality if and only if K and L are homothetic.

Taking for $p = n - i$ in (2.1.1), (2.1.1) changes to

$$S_{\tilde{w}_i}(I_j(K, L), I_j(D, D'))^{n-1} \leq S_{\tilde{w}_i}(IK, ID)^{n-j-1} S_{\tilde{w}_i}(IL, ID')^j,$$

with equality if and only if K and L are dilates.

This is just a new result which was given by Zhao and Leng [22].

Let D and D' be single point and taking for $p = n$ in (2.1.1), (2.1.1) changes to the well know Minkowski inequality for mixed projection bodies.

References

- [1] H. Busemann, A theorem on convex bodies of the Brunn-Minkowski type, *Proc. Nat. Acad. Sci. USA.*, **35**(1949), 27-31.
- [2] E. Lutwak, Intersection bodies and dual mixed volumes, *Adv. Math.*, **71**(1988), 232-261.
- [3] G. Zhang, Centered bodies and dual mixed volumes, *Trans. Amer. Math. Soc.*, **345**(1994), 777-801.
- [4] H. Busemann, The foundations of Minkowskian geometry, *Comment. Math. Helvet.*, **24**(1950), 156-186.
- [5] K. Ball, Shadows of convex bodies *Trans. Amer. Math. Soc.*, **327**(1991), 891-901.
- [6] K. Ball, Volume ratios and a reverse isoperimetric inequality, *J. London Math. Soc.*, **44**(1991), 351-359.
- [7] J. Bourgain, On the Busemann-Petty problem for perturbations of the ball, *Geom. Funct. Anal.*, **1**(1991), 1-13.
- [8] R. J. Gardner, Intersection bodies and the Busemann-Petty problem, *Trans. Amer. Math. Soc.*, **342**(1994), 435-445.
- [9] R. J. Gardner, A positive answer to be Busemann-Petty proplem in three dimensions, *Ann. Math.*, **140**(1994), 435-477.
- [10] R. J. Gardner, "Geometric Tomography," Cambridge: Cambridge University Press, 1995.

- [11] R. Schneider, "Convex bodies: The Brunn-Minkowski Theory", Cambridge: Cambridge University Press, 1993.
- [12] E. Lutwak, Centroid bodies and dual mixed volumes, *Proc. London Math. Soc.*, **60**(1990), 365-391.
- [13] E. Lutwak, Mixed projection inequalities, *Trans. Amer. Math. Soc.*, **287**(1985), 92-106.
- [14] E. Lutwak, Dual mixed volumes, *Pacific J. Math.*, **58**(1975), 531-538.
- [15] E. Lutwak, Inequalities for mixed projection, *Tans. Amer. Math. Soc.*, **339**(1993), 901-916.
- [16] E. Lutwak, On quermassintegrals of mixed projection bodies, *Geom. Dedicata*, **33**(1990), 51-58.
- [17] E. Lutwak, The Brunn-Minkowski-Firey Theory I: Mixed Volumes and the Minkowski proplem, *J. Diff. Geometry.*, **38**(1993), 131-150.
- [18] E. Lutwak, The Brunn-Minkowski-Firey Theory II: Affine and Geominimal surface area, *Adv. Math.*, **118**(1996), 244-294.
- [19] G. S. Leng, The Brunn-Minkowski inequaily for volume difference, *Adv. Appl. Math.*, **32**(2004), 615-624.
- [20] W. Süß, Zusammensetzung von Eikörpern und homothetische Eiflächen, *Tôhoku. Math. J.*, **35**(1932), 47-50.
- [21] G. H. Hardy, J. E. Littlewood and G. Pölya, "Inequalities", Cambridge Univ. Press. Cambridge, 1934.
- [22] C. J. Zhao, G. S. Leng, Inequalities for dual Quermassintegrals of mixed intersection bodies, *Proc. Indian Acad. Sci.*, **115**(2005), 79-91.

