

ON SEMILINEAR ELLIPTIC EIGENVALUE PROBLEMS

DAN PASCALI

The existence of eigenfunctions and eigenvalues of the semilinear equation given below is reduced to the study of a constrained critical point problem. For each $r \neq 0$, the corresponding functional admits two local minima while the third critical point is obtained by a mountain pass technique.

More precisely, given $r \in \mathbb{R}$, we look for eigensolutions $(\lambda, u) \in (\mathbb{R}, H_0^1(\Omega))$ of the semilinear equation of the form

$$\begin{cases} Lu = \lambda f(x, u) & \text{in } \Omega, \\ (Lu, u) = 2r, & r \neq 0, \end{cases} \quad (P)$$

where $\Omega \subset \mathbb{R}^N$ is a bounded domain with a smooth boundary $\partial\Omega$. The last relation normalizes the eigensolution $u(x)$. The second order linear operator

$(Lu)(x) = -\partial_i(a_{ij}(x)\partial_j u(x)) + c(x)u(x)$, $\partial_i = \frac{\partial}{\partial x_i}$, $i, j = 1, 2, \dots, N$, with the summation convention, is uniformly elliptic and selfadjoint ($a_{ij} = a_{ji}$ and $0 \leq c \in L^\infty(\Omega)$). Upon $f(x, t)$ we make the following hypotheses:

a) $f \in C(\bar{\Omega} \times \mathbb{R}, \mathbb{R})$, $f(x, t)t \geq 0$ and there is a neighborhood U of zero such that $f(x, t)t > 0$ for all $t \in U \setminus \{0\}$ and $x \in \Omega$;

b) $|f(x, t)| \leq k_1 + k_2|t|^p$ with $1 \leq p < \frac{N+2}{N-2}$ if $N \geq 3$ and p is unrestricted if $N = 1, 2$.

Define $G(u) = \frac{1}{2}(Lu, u)$, $\partial G_r = \{u \in H_0^1(\Omega) \mid G(u) = r\}$ as well as

$$F(x, u) = \int_0^u f(x, t)dt \quad \text{and} \quad J(u) = - \int_\Omega F(x, u(x))dx.$$

One can see as in [9, Appendix B] that

1°. $J \in C^1(H_0^1(\Omega), \mathbb{R})$, so J' is locally Lipschitz continuous;

2°. J' is completely continuous, i.e., it carries weakly convergent sequences into strongly convergent ones;

3°. J is weakly continuous in $H_0^1(\Omega)$.

In our setting ∂G_r is regular, i.e. $(G'u, v) \neq 0$ for any $v \in \partial G_r$. The tangent space to ∂G_r at u coincides with the null space $N(G'u)$ and so it has the form

$$T_u = T_u(\partial G_r) = \{ v \in H_0^1(\Omega) \mid (G'u, v) = 0 \}.$$

Recall that $u \in \partial G_r$ is said to be a *critical point of J constrained on ∂G_r* if $(J'(u), v) = 0$ for all $v \in T_u$. It is easily seen that any local extremum of the restriction $J|_{\partial G_r}$ is a critical point [6].

Introduce on ∂G_r the mapping

$$J'_r(u) = J'(u) - \frac{(J'(u), u)}{(G'u, u)} G'u.$$

We can show that $u \in \partial G_r$ is a constrained critical point of J to ∂G_r if and only if $J'_r(u) = 0$.

Indeed, define the linear projection from $H_0^1(\Omega)$ onto the tangent space T by setting

$$P_u v = v - \frac{(G'u, v)}{(G'u, u)} u \quad u \in H_0^1(\Omega)$$

and the above equivalence is a result of the relations

$$(J'_r(u), (I - P_u)v) = 0$$

$$(J'_r(u), P_u v) = ((J'(u), P_u v).$$

Therefore, a constrained critical point of J to ∂G_r corresponds to a weak solution of (P) with

$$\lambda = \frac{(G'u, u)}{(J'(u), u)}.$$

Therewith $c = J_r(u)$ is called a *critical value* of J_r .

REMARK 1. That $(J'(u), u) \neq 0$ follows from the hypothesis a); indeed, by [12, Lemma 1.0], if $u \neq 0$, then $\text{meas}\{x \in \Omega \mid |u(x)| \leq \varepsilon\} > 0$ for any $\varepsilon > 0$, and so $f(x, u(x)) u(x) > 0$ on a set of positive measure.

Hence, there is a one-to-one correspondence between the critical points of J_r and the weak solutions of the problem (P). If $f(x, \cdot)$ is locally Lipschitz continuous, a standard bootstrap argument shows that u is in fact a classical solution.

Several authors have studied the existence of multiple solutions of problem (P) under the additional restriction that $f(x, \cdot)$ is odd [1] while a result when $f(x, \cdot)$ is non-odd and ∂G_r sphere-like is given in [14]. A more general case is approached in [13].

The functional J is said to satisfy the Palais-Smale condition at level c , $c \in \mathbb{R}$ ((PS)_c for short), if every sequence $\{u_n\} \subset \partial G_r$ with $J_r(u_n) \rightarrow c$ and $J'_r(u_n) \rightarrow 0$ as $n \rightarrow \infty$ contains a convergent subsequence. Moreover, J_r satisfies (PS)₋ or (PS)₊ if it meets (PS)_c for all $c < 0$ or $c > 0$, respectively.

Similarly as in [12], we can prove

LEMMA. Let $\{u_n\} \subset \partial G_r$ be a sequence such that $J_r(u_n) \rightarrow c$ as $n \rightarrow \infty$. Then $\{u_n\}$ is bounded in $H^1_0(\Omega)$.

Consider on $H^1_0(\Omega)$ a more convenient inner product given by

$$\langle u, v \rangle = \int_{\Omega} (a_{ij}(x) \partial_i u \partial_j v + c(x) uv) dx$$

so that the corresponding norm $\|u\| = \langle u, u \rangle^{\frac{1}{2}}$ is equivalent to the initial norm, by virtue of the Poincaré inequality.

PROPOSITION [7]. The functional J_r satisfies (PS)_c for $c > 0$.

PROOF. If $J_r(u_n) \rightarrow c > 0$ and $J'_r(u_n) \rightarrow 0$, then $\{u_n\}$ is bounded according to the previous lemma and $u_n \rightharpoonup u_0$ in $H^1_0(\Omega)$, at least for a subsequence. Then $J_r(u_n) \rightarrow J_r(u_0) = c > 0$ by the weak continuity of J_r and $u_0 \neq 0$.

Denoting

$$\mu_n = \frac{(J'(u_n), u_n)}{(G'u_n, u_n)}$$

we have $J'_r(u_n) = J'(u_n) - \mu_n G'u_n \rightarrow 0$. If $\mu_n \rightarrow 0$ then $J'(u_n) \rightarrow J'(u_0) = 0$ contradicting $(J'(u_0), u_0) > 0$ in the REMARK 1. If $\mu_n \rightarrow \infty$ then $G'u_n \rightarrow 0$

contradicts $(G'u_n, u_n) = 2r \neq 0$. Therefore, by passing to a subsequence, $\mu_n \rightarrow \mu_0 \neq 0$ which easily implies the strong convergence of $\{G'u_n\}$. Since $(G'u_n, u_n) = \|u_n\|^2$, we infer that $\{u_n\}$ converges strongly. ■

We obtain a positive and a negative solution by cutting the nonlinearity above or below zero,

$$f^+(x, t) = \begin{cases} f(x, t) & \text{if } t \geq 0, \\ 0 & \text{if } t < 0, \end{cases} \quad \text{and} \quad f^-(x, t) = \begin{cases} 0 & \text{if } t > 0, \\ f(x, t) & \text{if } t \leq 0, \end{cases}$$

as well as

$$F^\pm(x, u) = \int_0^u f^\pm(x, t) dt \quad \text{and} \quad J^\pm(u) = - \int_\Omega F^\pm(x, u(x)) dx$$

and then minimizing the corresponding functionals $J^+(u)$, $J^-(u)$ over ∂G_r . Note, by a), that J^\pm are nonpositive on ∂G_r .

Let us first consider the minimization problem

$$\inf_{u \in \partial G_r} J^+(u) = \alpha, \quad \alpha < 0,$$

and select a minimizing sequence $\{u_n\} \subset \partial G_r$ so that $u_n \rightharpoonup u^+$ in $H_0^1(\Omega)$.

Due to the weak continuity we have $J^+(u^+) = \lim J^+(u_n)$ implying $-\infty < \alpha$ and $u^+ \neq 0$. One can show, by contradiction as in [11], that $G(u^+) = r$ and so J^+ achieves its minimum on ∂G_r at the point u^+ . By applying ultimately the Lagrange multiplier rule, there is some $\lambda \in \mathbb{R}$ such that u^+ is a weak solution in $H_0^1(\Omega)$ of the equation (P). Now, according to L^p -theory [4], $u^+ \in C^{1,\alpha}(\bar{\Omega})$, and via the maximum principle, we infer that

$$Lu^+ = \lambda f(x, u^+), \quad u^+(x) > 0 \quad \text{for } x \in \Omega.$$

It is readily seen that $\lambda > 0$ and, by Hopf's boundary point lemma,

$$\frac{\partial u^+}{\partial n}(x) < 0 \quad \text{for } x \in \partial\Omega.$$

We conclude that $J^+(u) = J(u)$ in a C^1 -neighborhood of u^+ and hence u^+ is a local minimizer in the C^1 -topology of J restricted to ∂G_r .

Replace now f^+ and J^+ by f^- and J^- and obtain again a nonpositive local minimizer u^- in the C^1 -topology of J restricted to ∂G_r .

Related to our problem, we are ultimately interested in minimizing of J on the topology of $H_0^1(\Omega)$ and clearly an H_0^1 -neighborhood is much larger than a C^1 -neighborhood. In overtaking this step, we apply now the following criterion of H.Brezis-L.Nirenberg [2] and generalized by H.T.Tehrani [11]:

Assume $u_0 \in H_0^1(\Omega)$ is a local minimizer of J in the C^1 - topology restricted to ∂G_r . If $(J'(u_0), G'u_0) < 0$, then u_0 is also a minimizer of J in the H_0^1 - topology restricted to ∂G_r .

THEOREM. Suppose that f satisfies hypotheses a), b). Then problem (P) has at least three solutions.

PROOF. Pass to the new inner product and have $G(u) = \|u\|^2$. To carry out the above criterion, we need to show that $(J(u^\pm), u^\pm) < 0$. For u^+ , say, one obtains

$$\lambda \int_{\Omega} f(x, u^+) u^+ dx = \int_{\Omega} (Lu^+, u^+) dx = 2r \neq 0$$

whence $\lambda > 0$, by assumption b), and hence

$$(J'(u^+), u^+) = - \int_{\Omega} f(x, u^+) u^+ dx = - \frac{1}{\lambda} < 0.$$

According to the Brezis-Nirenberg-Tehrani result, u^\pm are also local minimizers of J constrained to ∂G_r with H^1 - topology. Since the restriction of J on ∂G_r satisfies the $(PS)_+$ condition and J has two local minima according to a variant of the Mountain Pass Theorem (see e.g. [3, Corollary 2] or [10, Theorem 9.3]) there exists a third critical point of J on ∂G_r . ■

This theorem is essentially due to Li Yongqing [13] under more restrictive conditions upon the nonlinearity f :

a') $f \in C^1(\bar{\Omega} \times \mathbb{R}, \mathbb{R})$, $f(x, t)t \geq 0$ and $f(x, t) = 0$ iff $t = 0$;

b') $|\partial_t f(x, t)| \leq k_1 + |t|^{p-1}$ with $1 \leq p \leq \frac{N+2}{N-2}$ if $N \geq 3$ and $1 \leq p < \infty$ if $N = 1, 2$.

For the existence of the third solution, a retractable property of f is proved.

In the superlinear case, the constrained critical points under conditions a'), b') have been considered earlier (see, e.g. [5]).

REFERENCES

- [1] A.AMBROSETTI, Variational Methods and Nonlinear Problems: Classical results and recent advances, in Topological Nonlinear Analysis: Degree, Singularity and Variations, 1-36, (M.Matzeu, A.Vignoli, eds.), Birkhauser, Boston, 1995.
- [2] H.BREZIS and L.NIRENBERG, H^1 versus C^1 local minimizers, *C.R.Acad.Sci.Paris Ser A-B* 317 (1993), A465 - A473.
- [3] K.C.CHANG, On the mountain pass lemma, *Proc. Equadiff 6, Brno, Czechoslovakia*, 203-208, *Lect.Notes in Math.* 1192, Springer-Verlag, Berlin, 1986.
- [4] D.GILBARG and N.S.TRUDINGER, Elliptic Partial Differential Equations of Second Order, *Grundlehren* 224, Springer-Verlag, Berlin, 1977.
- [5] O.KAVIAN, Introduction a la theorie des points critiques, Springer-Verlag, Paris, 1993.
- [6] D.PASCALI, Nonlinear variational eigenvalue inclusions, in Differential Equations, Proc. EQUADIFF 87, Xanthi-Grece, (C.Defermos, ed.), *Lect.Notes Pure Appl.Math.Math.* 118, 543-551, Marcel Dekker Inc, New York, 1989.
- [7] D.PASCALI, Generalized subdifferentiable perturbations, *Proc.Intern.Conf. on Theory Appl.Diff.Equations*, vol.2, 277-281, (A.R.Aftabizadeh, ed.), Ohio Univ. Press, Athens, 1989.
- [8] D.PASCALI, Elliptic eigenvalue variational inequalities, *Libertas Math.* XII (1992), 39-56.
- [9] P.RABINOWITZ, Minimax Methods in Critical Point Theory with Applications to Differential Equations, C.B.M.S. 65, Amer.Math.Soc., Providence, 1986.
- [10] M.STRUWE, Variational Methods, Springer-Verlag, 1990.
- [11] H.T.TEHRANI, On some semilinear elliptic boundary value problems, Ph.D. thesis, Courant Inst., New York Univ., Sept.1994.
- [12] LI YONGQING, On a nonlinear elliptic eigenvalue problems, *J.Differential Equations* 117 (1995), 151-164.
- [13] LI YONGQING, Three solutions of a semilinear elliptic eigenvalue problem, Preprint 1993, Dept.of Math., Univ.of Stockholm.
- [14] E.ZEIDLER, Nonlinear functional analysis and its applications, Part III, Variational Methods and Optimization, Springer-Verlag, New York, 1985.
- [15] E.ZEIDLER, Applied Functional Analysis, Applications to Mathematical Physics, *Appl.Math.Sc.* 108, Springer-Verlag, New York, 1995.

ADDITIONAL REFERENCES

- H.AMANN, Lusternik-Schnirelman theory and non-linear eigenvalue problems, *Math. Ann.* 199 (1972), 55-72.
- A.AMBROSETTI, Critical points and nonlinear variational problems, *Mem. Soc. Math. France (N.S.)* 49(1992), 139p M. R. 93g 58024
- K.-C.CHANG, Infinite Dimensional Morse Theory and its Applications, *Sem. Math. Sup. 97, Les Presses de l'Univ. de Montreal*, 1985.
- K.- C.CHANG, Infinite Dimensional Morse Theory and Multiple Solution Problems, *Birkhauser, Boston*, 1993.
- D.G.COSTA and J.V.A.GONÇALVES, Critical point theory for nondifferentiable functionals and applications, *J. Math. Anal. Appl.* 153(1990), 470-485. M. R. 91j 58034
- I.EKELAND, Dual variational methods in non-convex optimization and differential equations, *Lect. Notes Math.* 979, 108-119, *Springer-Verlag, Berlin*, 1983.
- I.EKELAND and J.M.LASRY, Problemes variationnels non convexes en dualite, *C.R. Acad. Sci. Paris Ser A-B* 291(1980), no. 8, A493-A496. M. R. 82g 58021
- I.EKELAND and J.M.LASRY, Duality in nonconvex variational problems, *Advances in Hamiltonian systems (Rome, 1981)*, *Ann. CEREMADE, Birkhauser, Boston*, 1983. M. R. 85c 49011
- D.G. de FIGUEREIDO, Lectures on the Ekeland variational principle with applications and detours, *Tata Institute Fund. Res. Lect. in Math.* 81, *Bombay, Springer-Verlag*, 1989.
- S.S.LIN, On the number of positive solutions for nonlinear elliptic equation when a parameter is large, *Nonlinear Anal.* 16 (1991), 283-297.
- P.L.LIONS, On the existence of positive solutions of semilinear elliptic equations, *SIAM Rev.* 24 (1982), 441-467.
- J.MAWHIN and M.WILLEM, Critical Point Theory and Hamiltonian Systems, *Appl. Math. Sci.* 74, *Springer-Verlag, New York*, 1989.
- S.I.POKHOZHAEV, On the method of fibering a solution in nonlinear boundary value problems, *Proc. Steklov Inst. Math.* 192, 157-173, *Amer. Math. Soc*, 1992.
- P.RABINOWITZ, Critical point theory, in *Topological Nonlinear Analysis, Degree, Singularity and Variations*, 464-513, (*M. Matzeu, A. Vignoli, eds.*), *Birkhauser, Boston*, 1995.
- M.SCHECHTER and K.TINTAREV, Spherical maxima in Hilbert space and semilinear elliptic eigenvalue problems, *J. Diff. Int. Equations* 3 (1990), 889-899.
- M.SCHECHTER and K.TINTAREV, Pairs of critical points produced by linking subsets with applications to semilinear elliptic problems, *Bull. Soc. Math. Belg.* 44 (1992), 249-261.

A.SZULKIN, Minimax principles for lower semicontinuous functions and applications to nonlinear boundary value problems, *Ann.Inst.H.Poincaré, Analyse non linéaire* 3 (1986), 77-119.

K.TINTAREV, Level set maxima and quasilinear elliptic problems, *Pacific J. Math.* 153 (1992), 185-200. M. R. 93d:47109

Z.Q.WANG, On a superlinear elliptic equation, *Ann.Inst.H.Poincaré, Analyse non linéaire* 8 (1991), 43-58.

E.ZEIDLER, The Ljusternik-Schnirelman theory for indefinite and not necessarily odd nonlinear operators and its applications, *Nonlinear Anal.* 4 (1980), 451-489.

E.ZEIDLER, Ljusternik-Schnirelman theory on general level sets, *Math.Nachr.* 129 (1986), 235-259.

X.-P.ZHU and H.-S.ZHOU, Existence of multiple positive solutions of inhomogeneous semilinear elliptic problems in unbounded domains, *Proc.Royal Soc. Edinburg* 115A (1990), 301-318.